

## Molecular Detection and Antimicrobial Activity of Enterolysin a Endopeptidase among *Enterococcus faecalis* Strains

Zeinab Hatami\*<sup>1</sup>; Mitra Salehi<sup>1</sup>; Nader Mosavari<sup>2</sup>; Roya Razavi poor<sup>1</sup>; Marzie Eyn nam<sup>1</sup>; Mohammad Jaam shahriyari<sup>1</sup>

1-Department of Microbiology, North Tehran Branch, Islamic Azad University, Tehran, Iran

2-Razi Vaccine and Serum Research Institute, Karaj, Iran

hatami\_oct@yahoo.com

**Background & Objectives:** Enterococci are one of the gastro- intestinal tract commensal bacteria in humans and animals with ability to produce antimicrobial peptids. Enterolysin A is a large (calculated molecular weight of 34,501 Da), heatlabile bacteriocin and therefore fits the general characteristics of class IV bacteriocins. Target of present study is Molecular detection of enterolysin A gene among animal isolates of *Enterococcus faecalis* for their potential as biopreservatives in food or feed.

**Methods:** In this study occurrence of class IV enterocin structural gene (enterolysin A) in a target of *Enterococcus faecalis* isolated from 63 samples of different locales of Tehran animal faeces have been surveyed. Enterococcal strains were isolated from other faeces Gram-positive and negative bacteria using Bile Aesculin Azide Agar medium and after strains purifications, *E. faecalis* species identification and occurrence of enterolysin A gene was performed by PCR Methods. Cell-free neutralized supernatant (of selected gene positive strains) was used to test bacteriocin production and antimicrobial spectrum of endopeptidase was assayed by diffusion Methods (disk) on the gram-positive and negative indicators bacteria.

**Results:** 10 strains of *E. faecalis* were purified from 63 isolated samples. 6 strains of this 10 *E. faecalis* strains (60%) had enterolysin A gene that they inhibited growth of indicator bacteria such as clinical strain of *Staphylococcus epidermidis*, *Listeria monocytogenes* *Bacillus subtilis* and *Bacillus cereus*.

**Concolusion:** This strains produce an endopeptidase, enterolysin A encoded by *enl A*, which is homologous to other cell wall lytic enzymes such as lysostaphin and zoocin A. However, unlike these enzymes enterolysin A has a broad spectrum of activity and can lyse a wide range of Gram-positive bacteria.

**Keywords:** *Enterococcus faecalis*; PCR; Enterolysin A